

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12Q 1/68</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/24460</p>
<p>(21) International Application Number: PCT/US96/20650 (22) International Filing Date: 23 December 1996 (23.12.96) (30) Priority Data: 08/579,787 28 December 1995 (28.12.95) US (71) Applicant: TROPIX, INC. [US/US]; 47 Wiggins Avenue, Bedford, MA 01730-2314 (US). (72) Inventors: BRONSTEIN, Irena, Y.; 11 Ivanhoe Street, Newton, MA 02158 (US). FORTIN, John, J.; 163 West Main Street, Georgetown, MA 01833 (US). MARTIN, Chris, S.; 74 Chester Road, Belmont, MA 02178 (US). VOYTA, John, C.; 99 Maynard Farm Road, Sudbury, MA 01776 (US). (74) Agents: CONLIN, David, G. et al.; Dike, Bronstein, Roberts & Cushman, L.L.P., 130 Water Street, Boston, MA 02109 (US).</p>		<p>(43) International Publication Date: 10 July 1997 (10.07.97)</p> <p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published With international search report.</p>
<p>(54) Title: MULTIPLE REPORTER GENE ASSAY</p>		
<p>(57) Abstract</p> <p>A method of measuring the activity of at least two reporter gene products in an aliquot of a sample extract is disclosed. The activities of a first and second reporter enzyme are quantified by measuring the light signal produced by degradation of a first substrate by the first reporter enzyme and the light signal produced by the degradation of a second substrate by a second reporter enzyme. Both quantifications are sequentially performed on the same aliquot of sample extract.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

MULTIPLE REPORTER GENE ASSAY

FIELD OF THE INVENTION

This invention relates to methods of detecting transcriptional activity in a cell using two or more reporter genes.

5 BACKGROUND OF THE INVENTION

Reporter gene assays are useful in the study of gene regulatory elements because reporter gene activity, i.e., production of the reporter protein, is directly proportional to transcriptional activity of the regulatory elements of the gene. A reporter gene construct for use in these assays contains one or more gene
10 regulatory elements which are of interest, the minimal sequence requirements for transcription of a functional mRNA and the coding sequence for a reporter protein. Alam, J., et al, *Anal. Bioch.*, 188: 245-254 (1990). Analysis of constructs containing various deletions within the regulatory region enables mapping of regulatory sequences necessary for transcription and cell specific expression.

15

The reporter protein typically has a unique enzymatic activity or structure which enables it to be distinguished from other proteins present. The activity of the transcribed reporter protein, or quantification of the expressed protein, provides an indirect measurement of gene expression. Reporter gene assays enable
20 the identification of sequences and factors that control gene expression at the transcriptional level. Bronstein, I., et al, *BioTechniques*, Vol. 17, No. 1, p. 172 (1994). Other uses for reporter gene assays include: identification of sequences and factors that control genes at the translational level, study of mechanisms and factors that influence and alter gene expression levels and drug screening in cell-
25 based assays.

In single reporter gene assays with poor sensitivity it is difficult to distinguish negative results caused by the lack of expression or low level assay

- 2 -

sensitivity. This problem can be overcome with assays of greater sensitivity. Multiple gene assays are commonly used to provide controls for efficiency of transfection. In such assays, cells are transfected with a mixture of two separate plasmids, each having a different reporter gene. The expression of one reporter gene is controlled by different regulatory regions being studied while the other
5 reporter gene, acting as a control, is generally constitutively expressed by a standard promoter or enhancer. The activity of the experimental reporter enzyme is normalized to the activity of the control reporter enzyme.

10 In the known examples of assaying multiple reporter gene expression, a separate aliquot of the sample must be used in a separate assay to test for the activity of each enzyme. Alam, J., et al, *Anal. Bioch.*, 188: 245-254 (1990). The necessity of testing a separate portion for each enzyme decreases the precision of the assay and may introduce experimental errors into the results. Therefore, a
15 multiple reporter gene assay which is sequentially performed on the same aliquot of cell extract would simplify the assay procedure and minimize experimental errors. The use of multiple reporter genes can improve the efficiency of high throughput screening for drug discovery.

20 To provide relevant experimental information, reporter assays must be sensitive, thus enabling the detection of low levels of reporter protein in cell lines that transfect poorly. The sensitivity of a reporter gene assay is a function of the detection method as well as reporter mRNA and protein turnover, and endogenous (background) levels of the reporter activity.

25 Commonly used detection techniques use isotopic, colorimetric, fluorometric or luminescent enzyme substrates and immunoassay-based procedures with isotopic or color endpoints. Many of these systems, however, have disadvantages that limit their usefulness in these assays. For example, isotopic
30 substrates and immunoassays are limited by the cost, sensitivity and inconvenience of using radioisotopes. Fluorometric systems require external light sources which must be filtered to discriminate fluorescent signal, thereby limiting the sensitivity

- 3 -

and increasing complexity of the detection system. Furthermore, fluorescence from endogenous source can interfere with fluorometric measurements.

Colorimetric systems lack the sensitivity desired for sensitive reporter gene assays. Chemiluminescent and bioluminescent assays, on the other hand, have been found
5 to be more rapid and sensitive than colorimetric assays and fluorometric assays.

Jain, V.K. and Magrath, I.T., *Anal. Biochem.*, 199: 119-124 (1991). It would be, therefore, desirable to have a multiple reporter gene assay as aforesaid, which uses a luminescent detection system.

10 A number of genes are currently used as reporter genes including chloramphenicol acetyltransferase (CAT), secreted alkaline phosphatase, luciferase, β -galactosidase, β -glucuronidase, and human growth hormone, among others. Bronstein, I., et al, *Anal. Biochem.*, 219: 169-181 (1994). β -galactosidase and CAT are two of the most widely used reporter genes. See Alam, et al., 1990). β -
15 galactosidase detection is commonly performed with colorimetric substrates which lack sensitivity. Fluorescent substrates are also used to detect β -galactosidase, however, those assays also lacks sensitivity and are limited by background autofluorescence and signal quenching. The most widely used assay for CAT is radioisotopic, exhibits only moderate sensitivity, and suffers from a narrow
20 dynamic range. β -Glucuronidase (GUS) is a very widely used reporter gene in plant genetic research and to a lesser extent in mammalian cells. A common assay for GUS uses a fluorescent substrate, but is limited by background autofluorescence and signal quenching. Luciferase has become a more widely used reporter gene as it is quantitated using a very sensitive bioluminescent assay
25 utilizing the substrate, luciferin.

Sensitive chemiluminescent assays, not limited to reporter gene assays, have been described using dioxetane substrates. Bronstein, U.S. Patent No. 4,978,614, incorporated herein by reference. These dioxetane substrates emit visible light
30 following enzyme induced degradation. Enhancement of the chemiluminescent degradation of 1,2-dioxetanes by enhancer substances comprised of certain water soluble substances, such as globular proteins that have hydrophobic regions, has

- 4 -

been described. Voyta et al., U.S. Patent No. 5,145,772, incorporated herein by reference. These dioxetane substrates are also used in reporter gene assays for alkaline phosphatase, β -galactosidase, and β -glucuronidase for example. See e.g., Bronstein, I., et al, *Anal. Biochem.*, 219: 169-181 (1994) and citations therein.

5 However, no reporter gene assay using dioxetane substrates has been described in which the products of multiple reporter genes are sequentially quantitated in the same aliquot of cell extract.

Simple, rapid and highly sensitive, combined multiple reporter gene assays
10 to detect commonly used reporter genes which do not use radioisotopes or require external light sources are highly desirable.

It is desirable to have a multiple reporter gene assay in which the reagents enhance the light signal produced by the reporter enzymes. It is also important
15 that the signal from one reporter enzyme in a multiple reporter gene assay does not significantly interfere with the signal from the other reporter enzymes during measurement of their maximum light signal. It would be useful to have an assay which produces enhanced levels of light and therefore increases assay dynamic range and sensitivity and enables the use of a wide variety of instruments.

20

SUMMARY OF THE INVENTION

The method of the present invention provides a rapid, highly sensitive, non-isotopic method for sequentially detecting multiple reporter gene products in a single aliquot of cell extract. The method of the present invention comprises
25 quantifying the activity of a first reporter enzyme by measuring the light signal produced by degradation of a substrate by the first reporter enzyme, and quantifying the activity of a second reporter enzyme by measuring the light signal produced by degradation of a second substrate by the second reporter enzyme, wherein both quantifications are sequentially performed on the same aliquot of
30 sample extract. In one preferred embodiment, the presence of the first substrate enhances the light signal produced by degradation of a second substrate.

- 5 -

In preferred embodiments, the method of the present invention further comprises decreasing the light signal produced from the first reporter enzyme prior to quantifying the activity of the second reporter enzyme. Decreasing the signal from the first reporter enzyme preferably comprises substantially inactivating the first reporter enzyme or decreasing the amount of the first substrate.

In the methods of the present invention, substantially inactivating the first reporter enzyme comprises altering the pH of the aliquot, heating the aliquot to degrade the first enzyme or adding specific reagents to decrease activity of the first enzyme, such as alcohols such as isopropanol or ethanol, surfactants such as cetyl trimethyl ammonium bromide (CTAB) or substrate analogs.

In the methods of the present invention, decreasing the amount of the first substrate comprises adding an additional amount of the first enzyme sufficient to degrade the first substrate.

The reporter enzymes useful in the practice of this invention include luciferase, β -galactosidase, β -glucuronidase, alkaline phosphatase, and secreted placental alkaline phosphatase. Preferably, at least one of the first or second reporter enzymes is a hydrolytic enzyme capable of reacting with a dioxetane substrate. Useful hydrolytic enzymes include alkaline and acid phosphatases, esterases, decarboxylases, phospholipase D, β -xylosidase, β -D-fucosidase, thioglucosidase, β -D-galactosidase, α -D-galactosidase, α -D-glucosidase, β -D-glucosidase, α -D-mannosidase, β -D-mannosidase, β -D-fructofuranosidase, β -D-glucosiduronase, and trypsin. In one preferred embodiment, the first reporter enzyme is luciferase. In preferred embodiments, the second reporter enzyme comprises β -galactosidase. In one particularly preferred embodiment the first reporter enzyme is luciferase and the second reporter enzyme is β -galactosidase.

Preferably, at least one of the first or second substrates is a dioxetane. Dioxetane substrates useful in the present invention include: 3-(2'-spiroadamantane)-4-methoxy-4-(3"-phosphoryloxy)phenyl-1,2-dioxetane, disodium

- 6 -

salt (AMPPD), or disodium 3-(4-methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)-tricyclo-[3.3.1.1^{3,7}]decan]-4-yl]phenyl phosphate (CSPD), 3-(2'-spiroadamantane)-4-methoxy-4-(3"-β-D-galactopyranosyl)phenyl-1,2-dioxetane (AMPGD); 3-(4-methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1.1^{3,7}]decan]-4-yl-phenyl-β-D-galactopyranoside (Galacton™), and 5-chloro-3-(methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1^{3,7}]decan-4-yl-phenyl-β-D-galactopyranoside (Galacton-Plus).

In certain embodiments, the method further comprises adding a water soluble polymeric enhancer molecule to enhance the light signal produced by enzymatic degradation of the dioxetane. There can be one enhancer for each dioxetane or the same enhancer can be used for multiple substrates. Polymeric enhancers useful in the practice of the present invention include bovine serum albumin, human serum albumin or polymeric quaternary onium salts. The polymeric quaternary onium salts comprise poly vinylbenzyltrimethylammonium chloride (TMQ), polyvinyl benzyl tributyl ammonium chloride (TBQ), polyvinylbenzyltributylphosphonium chloride, polyvinylbenzyl benzyltrimethylammonium chloride (BDMQ) or polyvinyl tributyl sulfonium chloride. Other co-polymers, such as water soluble quaternary ammonium-phosphonium, ammonium-sulfonium and sulfonium-phosphonium polymers are also useful as enhancer molecules.

In other embodiments, these methods further comprise adding an accelerator solution prior to measuring the second enzyme activity. The accelerator solution is used to substantially inactivate the first reporter enzyme and simultaneously increase the light signal produced by the second enzyme. The accelerator solution comprises a water soluble polymeric enhancer molecule. Preferably the pH of the accelerator is from about 9 to about 12. When the accelerator is added to the aliquot of cell extract it alters the pH of the aliquot. For example, in a preferred embodiment, the accelerator solution is at a pH at about 10.8. When the accelerator is added to the aliquot it increases the pH of the aliquot from about 6.0 to greater than 9.0. The altered pH of the aliquot decreases

- 7 -

the activity of the first reporter enzyme and activates the production of signal from accumulated enzymatic product of the degradation of the second substrate. The polymeric enhancers described in the preceding paragraph are used in the accelerator solution.

5

The method of this invention also provides methods of quantifying the product of more than two reporter genes by measuring multiple enzyme activities in a single aliquot of cell extract. One such method comprises (a) quantifying the activity of a first reporter enzyme in an aliquot of the cell extract by measuring the light signal produced by degradation of a first substrate; (b) quantifying the activity of a second reporter enzyme in the aliquot of the cell extract by measuring the light signal produced by degradation of a second substrate; and (c) quantifying the activity of a third reporter enzyme in the aliquot of the cell extract by measuring the light signal produced by degradation of a third substrate, wherein all quantifications are sequentially performed on the same aliquot of sample extract. In this method, the first, second and third substrates are different and at least one of the substrates is a dioxetane. In certain embodiments, the method of measuring products of more than two reporter genes further comprises decreasing the activity of the reporter enzymes prior to quantifying the activity of the subsequent reporter enzyme. For example, prior to quantifying the activity of the second reporter enzyme, the light signal generated by the first substrate for the first reporter enzyme is decreased. This method further comprises the measurement of the activity of the second reporter enzyme and then further measuring the activity of the third reporter enzyme.

15
20
25

The invention also provides a system for detecting the products of more than one reporter gene in an aliquot of a sample extract, the system comprising: the reagents for quantifying each of two or more reporter enzymes; the substrates for each of the reporter enzymes, wherein at least one of the substrates is a dioxetane; and an accelerator solution containing a water soluble polymeric enhancer molecule. The accelerator solution comprises a water soluble polymeric enhancer molecule at a pH from about 9 to about 12, to induce the signal produced

30

- 8 -

from the second or a third reporter enzyme-substrate reaction and decrease the activity of the first reporter enzyme. The polymeric enhancer comprises bovine serum albumin, human serum albumin or polymeric quaternary ammonium, sulfonium and phosphonium salts. The polymeric quaternary ammonium, sulfonium and phosphonium salts comprise polyvinylbenzyltrimethylammonium chloride (TMQ), polyvinylbenzyltributylammonium chloride (TBQ), polyvinylbenzyl benzyldimethylammonium chloride (BDMQ) polyvinylbenzylsulfonium chloride or polyvinylbenzyl tributylphosphonium chloride. Other co-polymers, such as ammonium-phosphonium, ammonium-sulfonium and sulfonium-phosphonium quaternary polymers are also useful as enhancer molecules.

One preferred embodiment of the invention provides a method of measuring the products of more than one reporter gene in an aliquot of a sample extract. The method comprises: (a) adding a first substrate which is the substrate of a first reporter enzyme product and a second substrate which is the substrate of a second reporter enzyme product to an aliquot of the cell extract, the first substrate comprising luciferin for detection of the first enzyme luciferase and the second substrate comprising a dioxetane and the second enzyme being a hydrolytic enzyme; (b) measuring the activity of the first reporter enzyme; (c) adding an accelerator solution which substantially inactivates the first reporter enzyme and simultaneously increases the chemiluminescent signal produced from the degraded substrate for the second reporter enzyme by increasing the pH of the aliquot, and (d) measuring the chemiluminescent signal produced from the degraded substrate for the second reporter enzyme in the same aliquot of the cell extract, wherein the presence of the first substrate enhances the light signal produced by degradation of a second substrate. In one preferred embodiment, the first reporter enzyme is luciferase and the second reporter enzyme is β -galactosidase.

The present invention provides a method of detecting the activity of a first and second reporter enzyme sequentially in the same aliquot of cell extract sample rather than using separate aliquots of the cell extract to individually measure each

- 9 -

enzyme as has been previously performed in the art. The method of the present invention therefore decreases the likelihood of experimental error, thus enabling more reliable data.

5 DETAILED DESCRIPTION OF THE INVENTION

The method of the present invention provides a rapid, highly sensitive, non-isotopic method for sequentially detecting multiple reporter gene activity in a single aliquot of cell extract sample. The method of the present invention is particularly useful in measuring transcriptional activity of cells transfected with more than one
10 reporter gene. Transfection of these cells is accomplished by methods known in the art. See e.g., Alam, J. and Cook, J.L., *Anal. Bioch.* 188: 245-254 (1990). Typically, cells are co-transfected with a DNA mixture of two separate plasmids, each having a different reporter gene. One plasmid has a reporter gene the expression of which is regulated by a known control promoter. This reporter gene
15 acts as a control. The second plasmid has a second reporter gene the expression of which is controlled by the regulatory region being studied. Transcription of each reporter gene is analyzed by measuring the activity of its product, a "reporter enzyme". The activity of the second reporter enzyme is typically normalized to the activity of the first reporter enzyme. *Id* at 249.

20

In one embodiment, the method of the present invention for measuring the activity of more than one reporter gene product in an aliquot of sample extract comprises the steps of: (a) quantifying the activity of a first reporter enzyme by measuring the light signal produced from degradation of a first substrate by the
25 first reporter enzyme; and (b) quantifying the activity of a second reporter enzyme by measuring the light signal produced from degradation of a second substrate by a second enzyme, wherein both quantifications are sequentially performed on the same aliquot of sample extract. The intensity of the luminescent signal produced from the substrates is a function of the activity of the enzyme, i.e., the
30 effectiveness of the enzyme in terms of its ability to degrade its substrate.

- 10 -


The first and second reporter enzymes useful in the present invention comprise any protein produced from any gene that exhibits enzymatic activity and degrades a substrate to produce a light signal. Examples of such enzymes include luciferase, alkaline phosphatase, β -galactosidase, β -glucuronidase, 5 carboxylesterase, lipases, phospholipases, sulphatases, ureases, peptidases, proteases and others. Further, in preferred embodiments, at least one of the reporter enzymes is a hydrolytic enzyme. In some of these embodiments, it is preferred that the second reporter enzyme is the hydrolytic enzyme. In other 10 embodiments, both enzymes are hydrolytic and can be selected from alkaline and acid phosphatases, esterases, decarboxylases, phospholipase D, β -xylosidase, β -D-fucosidase, thioglucosidase, β -D-galactosidase, α -D-galactosidase, α -D-glucosidase, β -D-glucosidase, β -D-glucuronidase, α -D-mannosidase, β -D-mannosidase, β -D-fructofuranosidase, β -D-glucosiduronase, and trypsin.

15 In the method of the present invention, the substrates for the reporter enzymes comprise any luminescent substrate capable of producing a light signal. Preferably, the first substrate is different from the second substrate and at least one substrate is a dioxetane. Dioxetanes are molecules that have a 4-membered ring in which 2 of the members are adjacent oxygen atoms. Dioxetanes can be thermally, 20 chemically or photochemically decomposed to form carbonyl products, e.g., esters, ketones or aldehydes. Release of energy in the form of light (i.e., luminescence) accompanies the decompositions. The dioxetane substrate contains an enzyme-cleavable group which can be cleaved by a corresponding enzyme. The reporter enzyme cleaves this enzyme-cleavable group from the substrate to form a 25 negatively charged group (e.g., an oxygen anion) bonded to the dioxetane. This dioxetane anion decomposes to form a luminescent substance which produces light. The light signal is detected as an indication of the presence and the amount of the enzyme. By measuring the intensity of the luminescence, the concentration of the reporter enzyme can be determined.

30

In one preferred embodiment, the dioxetane-containing substrate preferably has the formula

- 11 -



5

(I)

where T is a substituted (i.e. containing one or more C₁-C₇ alkyl groups or
10 heteroatom groups, e.g. halogens) or unsubstituted cycloalkyl ring (having between
6 and 12 carbon atoms, inclusive, in the ring) or polycycloalkyl group (having 2 or
more fused rings, each ring independently having between 5 and 12 carbon atoms,
inclusive), bonded to the 4-membered dioxetane ring by a spiro linkage, e.g., a
chloroadamantyl or an adamantyl group, most preferably chloroadamantyl; Y is a
15 fluorescent chromophore, (i.e. Y is group capable of absorbing energy to form an
excited, i.e. higher energy, state, from which it emits light to return to its original
energy state); X is hydrogen, a straight or branched chain alkyl or heteroalkyl
group (having between 1 and 7 carbon atoms, inclusive, e.g., methoxy,
trifluoromethoxy, hydroxyethyl, trifluoroethoxy or hydroxypropyl), an aryl group
20 (having at least 1 ring e.g., phenyl), a heteroaryl group (having at least 1 ring
e.g., pyrrolyl or pyrazolyl), a heteroalkyl group (having between 2 and 7 carbon
atoms, inclusive, in the ring, e.g., dioxane), an aralkyl group (having at least 1
ring e.g., benzyl), an alkaryl group (having at least 1 ring e.g., tolyl), or an
enzyme-cleavable group i.e., a group having a moiety which can be cleaved by an
25 enzyme to yield an electron-rich group bonded to the dioxetane, e.g., phosphate,
where a phosphorus-oxygen bond can be cleaved by an enzyme, e.g., acid
phosphatase or alkaline phosphatase, to yield a negatively charged oxygen bonded
to the dioxetane or OR; and Z is hydrogen, hydroxyl, or an enzyme-cleavable
group (as defined above), provided that at least one of X or Z must be an enzyme-
30 cleavable group, so that the reporter enzyme cleaves the enzyme-cleavable group
which leads to the formation of a negatively charged group (e.g., an oxygen
anion) bonded to the dioxetane, the negatively charged group causing the dioxetane

- 12 -

to decompose to form a luminescing substance (i.e., a substance that emits energy in the form of light) that includes group Y. The luminescent signal is detected as an indication of the activity of the reporter enzyme. By measuring the intensity of luminescence, the activity of the reporter enzyme can be determined.

5

When X, in formula I, is OR, moiety R is a straight or branched alkyl, aryl, cycloalkyl or arylalkyl of 1-20 carbon atoms. R may include 1 or 2 heteroatoms which may be P, N, S or O. The substituent R is halogenated. The degree of halogenation will vary depending on the selection of substituents on the adamantyl group, on the aryl group, and the desired enzyme kinetics for the particular application envisioned. Most preferably, R is a tribaloalkyl moiety. Preferred groups include trihalo lower alkyls, including trifluoroethyl, trifluoropropyl, heptafluoro butyrol, hexafluoro-2-propyl, α -trifluoromethyl benzyl, α -trifluoromethyl ethyl and difluorochloro butyl moieties. The carbon atoms of substituent R may be partially or fully substituted with halogens. When R is aryl, preferred groups may include a phenyl ring substituted with one or more chloro, fluoro, or trifluoromethyl groups, e.g., 2,5-dichlorophenyl, 2,4-difluorophenyl, 2,3,5-trifluorophenyl, 2-chloro-4-fluorophenyl or 3-trifluoromethyl phenyl. Fluorine and chlorine are particularly preferred substituents, although bromine and iodine may be employed in special circumstances.

15
20

Group Y is a fluorescent chromophore or fluorophore bonded to enzyme-cleavable group Z. Y becomes luminescent upon the dioxetane decomposition when the reporter enzyme cleaves group Z, thereby creating an electron-rich moiety which destabilizes the dioxetane, causing the dioxetane to decompose. Decomposition produces two individual carbonyl compounds, one of which contains group T, and the other of which contains groups X and Y. The energy released from dioxetane decomposition causes compounds containing the X and the Y groups to luminesce (if group X is hydrogen, an aldehyde is produced). Y preferably is phenyl or aryl. The aryl moiety bears group Z, as in formula I, and additionally 1-3 electron active groups, such as chlorine or methoxy, as described in U.S. Patent Application 08/231,673 filed April 25, 1994.

25
30

- 13 -

Any chromophore can be used as Y. In general, it is desirable to use a chromophore which maximizes the quantum yield in order to increase sensitivity. Therefore, Y usually contains aromatic groups. Examples of suitable chromophores are further detailed in U.S. Patent No. 4,978,614.

5

Group Z bonded to chromophore Y is an enzyme cleavable group. Upon contact with an enzyme, the enzyme-cleavable group is cleaved yielding an electron-rich moiety bonded to a chromophore Y; this moiety initiates the decomposition of the dioxetane into two individual carbonyl containing compounds e.g., into a ketone or an ester and an aldehyde if group X is hydrogen. Examples of electron-rich moieties include oxygen, sulfur, and amine or amino anions. The most preferred moiety is an oxygen anion. Examples of suitable Z groups, and the enzymes specific to these groups are given in Table 1 of U.S. Patent No. 4,978,614. Such enzymes include alkaline and acid phosphatases, esterases, 10 decarboxylases, phospholipase D, β -xylosidase, β -D-fucosidase, thioglucosidase, β -D-galactosidase, α -D-galactosidase, α -D-glucosidase, β -D-glucosidase, β -D-glucouronidase α -D-mannosidase, β -D-mannosidase, β -D-fructofuranosidase, β -D-glucosiduronase, and trypsin. 15

Dioxetane may also contain one or more solubilizing substituents attached to any of the T, Y and X, i.e., substituents which enhance the solubility of the dioxetane in aqueous solution. Examples of solubilizing substituents include carboxylic acids, e.g., acetic acid; sulfonic acids, e.g., methanesulfonic acid; and quaternary amino salts, e.g., ammonium bromide; the most preferred solubilizing 20 substituent is methane or ethanesulfonic acid. 25

Other dioxetanes useful in the practice of this invention are described in U.S. Patent No. 5,089,630; U.S. Patent No. 5,112,960; U.S. Patent Application 057,903 filed May 7, 1993 and U.S. Patent Application 08/231,673 filed April 25, 30 1994.

For assays using alkaline phosphatase or secreted human placental alkaline phosphatase (SEAP) as a reporter enzyme, the substrate for that enzyme preferably comprises a phosphate-containing dioxetane, such as 3-(2'-spiroadamantane)-4-methoxy-4-(3"-phosphoryloxy)phenyl-1,2-dioxetane, disodium salt (AMPPD), or
5 disodium 3-(4-methoxyspiro[1,2-dioxetane-3,2'(5'-chloro)-tricyclo-[3.3.1.1^{3,7}]decan]-4-yl]phenyl phosphate (CSPD), respectively.

For assays using β -galactosidase as a reporter enzyme, the substrate preferably comprises a dioxetane containing galactosidase-cleavable or
10 galactopyranoside groups. When using β -galactosidase, the luminescence results from the enzymatic cleavage of the sugar moiety from the dioxetane substrate. An example of such a substrate is 3-(2'-spiroadamantane)-4-methoxy-4-(3"- β -D-galactopyranosyl)phenyl-1,2-dioxetane (AMPGD), 3-(4-methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1.1^{3,7}]decan]-4-yl-phenyl- β -D-
15 galactopyranoside (Galacton™), and 5-chloro-3-(methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1^{3,7}]decan-4-yl-phenyl- β -D-galactopyranoside (Galacton-Plus™).

For assays using β -glucuronidase as a reporter enzyme, the substrate
20 comprises a dioxetane containing β -glucuronidase-cleavable groups such as a glucuronide. For assays using carboxyl esterase, the enzyme cleaves the ester group of the dioxetane. For assays using proteases and phospholipases, the enzymes cleave a suitable enzyme-cleavable group bound to the dioxetane.

25 In preferred embodiments of the assay of the present invention, the first and second substrates are both present in the aliquot during quantification of the activity of the first reporter enzyme. In these embodiments, the presence of the second reporter enzyme, as well as its substrate, does not interfere with the activity of the first reporter enzyme or measurement of the light signal produced
30 by degradation of the first substrate by the first reporter enzyme. In one preferred embodiment, light produced by degradation of the first substrate occurs at a pH at which the product of the degradation of the second substrate does not produce

- 15 -

light, or the second enzyme is not active during the first measurement. In one such embodiment, the product of enzymatic reaction with the second substrate does not generate a measurable signal during the first quantification because the environment of the aliquot inhibits light production by the product of this second enzymatic reaction. For example, the measurement of light production by the degradation of the first substrate preferably occurs at a neutral pH. The second reporter enzyme also acts on its substrate during this time, however, at this pH, no significant production of light from this reaction occurs. It is preferable to allow a predetermined period of time to elapse prior to increasing the pH and measuring light production of the second enzymatic reaction to allow the product of that reaction to accumulate, thereby producing a more intense light signal.

In other embodiments of the present invention, the presence of the second reporter enzyme in the aliquot does not interfere with the activity of the first reporter enzyme or measurement of the light signal produced by that first enzyme. In these embodiments, the method further comprises increasing the enzymatic activity of the second reporter enzyme subsequent to quantifying the activity of the first enzyme. This can be readily accomplished by methods known in the art to activate enzyme activity. In especially preferred embodiments, the pH of the aliquot is adjusted to create an environment in which the second enzyme is active prior to the second quantification. For example, in some embodiments, using alkaline phosphatase as the second reporter enzyme, the pH of the aliquot is raised because, the activity of alkaline phosphatase is increased at alkaline pH.

In other embodiments, the presence of the second enzyme does not interfere with the measurement of the light signal from the first substrate, because the substrate of the second enzyme is absent from the aliquot during this measurement. In such embodiments, the second enzyme is active but there is no substrate on which it can act. In one preferred embodiment, the activity of the second reporter enzyme is induced by the addition of the second substrate after decreasing the activity of the first reporter enzyme or after reduction in the signal produced by the first enzymatic reaction. For example, in an assay using β -galactosidase as the

- 16 -

first reporter enzyme and alkaline phosphatase as the second reporter enzyme, light production as a result of the β -galactosidase reaction is optimized at a higher than neutral pH. However, the increased pH simultaneously activates the second reporter enzyme, alkaline phosphatase, which could produce light which would interfere with the quantification of the activity of the first reporter enzyme. In such an assay, it is preferred that the second substrate is added subsequent to the first quantification and most preferably, after that light signal has abated. This addition is readily accomplished and is especially easy when using a detection device which has an injector system as is known in the art.

10

The light signal produced by the degradation of the first substrate by the first reporter enzyme does not interfere with the quantification of the light signal produced by the second reporter enzyme. This is accomplished in preferred embodiments by decreasing the signal produced by the substrate for the first reporter enzyme after measuring the light signal produced by degradation of the first substrate and prior to quantifying the activity of the second reporter enzyme.

15

In some preferred embodiments, the signal produced from the substrate for the first reporter enzyme is decreased by substantially inactivating the first reporter enzyme. Methods of substantially inactivating the first reporter enzyme are known in the art. However, these methods must not interfere with measurement of the second enzyme product activity. In one preferred method of this invention, the first reporter enzyme is substantially inactivated by altering the pH of the aliquot. The pH can be altered by adding acids or bases to the aliquot, depending on the enzyme, to provide an environment which is inhospitable to the first enzyme. Preferably the pH of the aliquot is increased. This inactivates the first enzyme, preventing it from further degrading its substrate and producing a light signal. In another embodiment, the aliquot is heated to degrade the first enzyme or specific inhibitors can be added. Examples of inhibitors include alcohols such as isopropanol or ethanol, surfactants such as cetyl trimethyl ammonium bromide (CTAB) or substrate analogs which bind to, and inactivate the enzyme.

20

25

30

- 17 -

In other embodiments, the signal produced by the substrate for the first enzyme is decreased by decreasing the amount of the first substrate. The appropriate method can be chosen by one skilled in the art based upon the specific enzyme and substrate used. However, in one such embodiment, the substrate concentration is preferably decreased by adding an additional amount of the first enzyme sufficient to degrade the residual first substrate remaining after quantification of the first reporter enzyme. For example, after measuring the light signal produced from the degradation of the first substrate by the first enzyme, e.g., alkaline phosphatase, an amount of alkaline phosphatase is added to the aliquot sufficient to degrade any residual substrate and to prevent any potentially interfering light signal. Since the second substrate is unique to the second reporter enzyme, the presence of the first enzyme in excess will not interfere with the detection of the second light signal. This is useful when dioxetane substrates are used for both the first and second reporter enzymes.

15

In another embodiment, decreasing the amount of the first substrate comprises heating the aliquot to degrade the first substrate. Heat inactivation is preferably utilized in embodiments in which the second substrate is added subsequent to inactivation of the first substrate.

20

In certain embodiments of the present invention, the presence of the first substrate enhances the light signal produced by degradation of the second substrate. This enhanced signal provides a more sensitive assay. Although the inventors do not intend to be bound by theory, it is believed that the energy produced by the degradation of the second substrate is transferred to the remaining first substrate, thereby enhancing the intensity of the light signal produced. In one preferred embodiment of this method, the first substrate is luciferin. In this embodiment the second substrate is preferably a dioxetane substrate. It is believed that luciferin is a very efficient energy acceptor and emits a light signal as a result of energy transfer from the excited state generated from the dioxetane decomposition. This produces greater signal intensity. For example, an enhanced signal is produced by 5-chloro-3-(methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1^{3,7}]decan-4-yl-

30

phenyl- β -D-galactopyranoside reaction with β -galactosidase in the presence of luciferin compared to the same reaction in the absence of luciferin. This enhanced signal provides a highly sensitive assay enabling the detection of β -galactosidase in the 8 fg to 20 ng range.

5

Certain preferred embodiments of the present invention further comprise adding a water soluble enhancer molecule which enhances the light signal produced from the enzymatic degradation of dioxetane substrates by the reporter enzymes.

10

Certain water soluble naturally-occurring and synthetic substances, generally macromolecular in nature, enhance the chemiluminescent signal intensity in part by providing a hydrophobic environment. These substances, for example water soluble globular proteins that contain hydrophobic regions: mammalian serum albumins such as bovine serum albumin (BSA) and human serum albumin

15

(HSA), or water soluble polymeric quaternary onium salts:

polyvinylbenzyltrimethylammonium chloride (TMQ) or

polyvinylbenzylbenzyltrimethylammonium chloride (BDMQ), and

polyvinylbenzyltributyl phosphonium chloride increase the chemiluminescent signal intensity produced by the decomposition of enzymatically cleavable 1,2-dioxetanes in aqueous solutions. Co-polymers, such as water soluble quaternary ammonium-phosphonium, ammonium-sulfonium and sulfonium-phosphonium polymers are useful as enhancer molecules.

By virtue of the presence of effective amounts of an enhancer substance or substances the intensity of the light emitted in aqueous medium is increased significantly as compared to the intensity of light emitted in the absence of such enhancers. These compounds enhance the intensity of the chemiluminescent signal from 1,2-dioxetanes by a factor of at least 10%, but usually at least tenfold and oftentimes by factors of at least 100 to 1,000,000.

30

Included among such enhancer substances are macromolecular globular proteins, generally ones having molecular weights ranging from about 1000 to


- 19 -

about 600,000 daltons, as determined by SDS gel electrophoresis, that include hydrophobic regions: mammalian serum albumins such as BSA, HSA and the like; globular proteins such as mammalian IgG, IgE, Protein A, avidins, and the like; serum lipoproteins, apolipoproteins, and the like.

5

Synthetic oligomeric or polymeric enhancer substances that can be used in practicing this invention include, first of all, water soluble polyvinylaryl quaternary onium salts, such as the polyvinylbenzyl quaternary ammonium, sulfonium and phosphonium salts having the formula:

10



15

20

(VI)

where E⁺ can be P, N or S.

In this formula each of R₁, R₂ and R₃ can be a straight or branched chain unsubstituted alkyl group having from 1 to 20 carbon atoms, inclusive, e.g., benzyl, methyl, ethyl, n-butyl, t-butyl, cetyl, or the like; a straight or branched chain alkyl group having from 1 to 20 carbon atoms, inclusive, substituted with one or more hydroxy, alkoxy, e.g., methoxy, ethoxy, benzyloxy or polyoxethylethoxy, aryloxy, e.g., phenoxy, amino or substituted amino, e.g., methylamino, amino, e.g., acetamido or cholesteryloxycarbonylamido, or fluoroalkane or fluoroaryl, e.g., heptafluorobutyl, groups, an unsubstituted monocycloalkyl group having from 3 to 12 ring carbon atoms, inclusive, e.g., cyclohexyl or cyclooctyl, a substituted monocycloalkyl group having from 3 to 12 ring carbon atoms, inclusive, substituted with one or more alkyl, alkoxy or fused benzo groups, e.g., methoxycyclohexyl or 1,2,3,4-tetrahydronaphthyl, a

35

- 20 -

polycycloalkyl group having 2 or more fused rings, each having from 5 to 12 carbon atoms, inclusive, unsubstituted or substituted with one or more alkyl, alkoxy or aryl groups, e.g., 1-adamantyl or 3-phenyl-1-adamantyl, an aryl, alkaryl or aralkyl group having at least one ring and from 6 to 20 carbon atoms *in toto*,
5 unsubstituted or substituted with one or more alkyl, aryl, or fluoroalkane or fluoroaryl groups, e.g., phenyl, naphthyl, pentafluorophenyl, ethylphenyl, benzyl, hydroxybenzyl, phenylbenzyl or dehydroabietyl; at least two of R₁, R₂ and R₃, together with the quaternary atom to which they are bonded, can form a saturated or unsaturated, unsubstituted or substituted nitrogen-containing, nitrogen and
10 oxygen-containing or nitrogen and sulfur-containing ring having from 3 to 5 carbon atoms, inclusive, and 1 to 3 heteroatoms, inclusive, and which may be benzoanlyated, e.g., 1-pyridyl, 1-(3-alkyl or aralkyl)imidazolium, morpholino, piperidino or acylpiperidino, benzoxazole, benzthiazole or benzamidazole.

15 The symbol X⁻ represents a counterion which can include, alone or in combination, moieties such as halide, i.e., fluoride, chloride, bromide or iodide, sulfate, alkylsulfonate, e.g., methylsulfonate, arylsulfonate, e.g., p-toluenesulfonate, substituted arylsulfonate, e.g., anilino-naphthylsulfonate (various isomers), lucifer yellow CH and diphenylanthracenesulfonate, perchlorate,
20 alkanolate, e.g., acetate, arylcarboxylate, e.g., fluorescein or fluorescein derivatives, benzoheterocyclic arylcarboxylate, e.g., 7-diethylamino-4-cyanocoumarin-3-carboxylate, or substituted monoaryloxyphosphate, e.g., a 3-(2'-spiroadamantane)-4-methoxy-(3"-phosphoryloxy)phenyl-1,2-dioxetane dianion or other dianions indicated in formula (I), *supra*.

25


The symbol n represents a number such that the molecular weight of such poly vinylbenzyl quaternary onium salts will range from about 800 to about 200,000, and preferably from about 20,000 to about 70,000, as determined by intrinsic viscosity or LALLS techniques.

30

Illustrative of such water soluble poly(vinylbenzyl quaternary ammonium salts) are TMQ, BDMQ, and the like.

- 21 -

Water soluble acetals of a polyvinylalcohol and a formylbenzyl quaternary onium salt, having the formula:



10

(VII)

wherein E⁺ can be P, N or S and each R₄ is the same or a different aliphatic substituent and X₁ is an anion, as disclosed and claimed in Bronstein-Bonte et al. U.S. Pat. No. 4,124,388 can also be used as enhancer substances in practicing this invention. And, the individual vinylbenzyl quaternary onium salt monomers used to prepare the poly vinylbenzyl quaternary onium salts of formula II above can also be copolymerized with other vinylbenzyl quaternary onium salt monomers whose polymers are depicted in formula II, or with other ethylenically unsaturated monomers having no quaternary onium functionality, to give polymers such as those disclosed and claimed in Land et al. U.S. Pat. No. 4,322,489; Bronstein-Bonte et al. U.S. Pat. No. 4,340,522, Land et al. U.S. Pat. No. 4,424,326; Bronstein-Bonte et al. U.S. Pat. No. 4,503,138 and Bronstein-Bonte U.S. Pat. No. 4,563,411, all of which polymers can also be used as enhancer substances in practicing this invention. Preferably these quaternized polymers will have molecular weights within the ranges given above for the polyvinylbenzyl quaternary ammonium salts of formula II.

Other water soluble oligomeric, homopolymeric and copolymeric materials can be used as enhancer substances in addition to or instead of the foregoing polymers and are further described in U.S. Pat. No. 5,145,772.

- 22 -

The amount of enhancer substance used varies depending on the particular enhancer chosen and the amount and type of chemiluminescent compounds present. The necessary amount can be readily determined by one skilled in the art based upon the present teachings. Furthermore, the disclosure contained within U.S. Pat. No. 5,145,772 would assist one skilled in the art in practicing this invention.

The enhancer molecule can be added at any point during the method of the present invention. If the first substrate, but not the second, is a dioxetane, the enhancer is preferably added prior to or simultaneous to quantification of the first reporter enzyme. If the second substrate, but not the first, is a dioxetane, the enhancer is added at any time, but preferably subsequent to quantification of the first reporter enzyme and prior or simultaneous to quantification of the second reporter enzyme. If both the first and second substrates are dioxetanes, the enhancer molecule is preferably added prior to any quantification or simultaneous to the first. In this last embodiment, one enhancer can be used to enhance the light signals produced by decomposition of both substrates. Or alternatively, a different enhancer is used for each substrate.

In some embodiments there is an incubation period subsequent to quantification of the light signal produced by the substrate for the first enzymes, in part, to allow for the light signal produced by the first substrate to diminish. The length of the incubation period will depend on the concentration of the first reporter enzyme and the half-life of the light signal produced by the substrate. If the first reporter is present in high concentrations, it may interfere with the quantification of low levels of the subsequent reporter enzymes. A longer incubation period will decrease the residual light signal from the substrate for the first reporter enzyme. The incubation period is preferably 10 to 120 minutes, more preferably 20 to 90 minutes, and most preferably 30 to 60 minutes. Taking into account the teachings of the present invention the appropriate length of the incubation period can readily be determined by one skilled in the art. Regardless of the incubation time period selected, the timing of the addition of the accelerator

- 23 -

and measurement of the signal produced by the second reporter enzyme after addition of the accelerator must be held constant throughout the assay.

The present invention provides a method of measuring the activity of more than two reporter gene products. In one preferred embodiment, this method
5 comprises first quantifying the activity of a first reporter enzyme in an aliquot of the cell extract by measuring the light signal produced by degradation of a first substrate and then decreasing the activity of the first reporter enzyme. The method also comprises quantifying the activity of a second reporter enzyme in the aliquot
10 of the cell extract by measuring the light signal produced by degradation of a second substrate. The method further comprises quantifying the activity of a third reporter enzyme in the aliquot of the cell extract by measuring the light signal produced by degradation of a third substrate. All quantifications are sequentially performed on the same aliquot of sample extract.

15

The method of the present invention may be performed in any luminometer, preferably luminometers with automatic injectors, or other instrumentation which enables the measurement of light emission. The method is also useful in a luminometer equipped with a single injector. However, when using a single
20 injector it is preferred that the injector is rinsed thoroughly between injection of the second buffer and the accelerator. The method can be performed using manual injection if the light signal of each sample is measured after approximately the same time interval subsequent to the addition of the accelerator to the samples.

25

A number of factors determine the volume of the aliquot of cell extract used, such as, the availability of the sample, instrumentation and availability of reagent. Preferably, the volume of the aliquot of cell extract used ranges from 1 to 100 μL , more preferably 2 to 20 μL , and most preferably from about 2 to 10 μL . The lysis solution used to lyse transfected cells to obtain the cell extract can
30 be any lysis solution known in the art.

- 24 -

The method of this invention is highly sensitive and detects from about 1 fg to 20 ng of purified luciferase, 10 fg to 20 ng of purified β -galactosidase, 2 fg to 100 ng of purified alkaline phosphatase, and 10 fg to 100 ng of purified β -glucuronidase.

5

In one preferred embodiment, the activity of more than one reporter gene product in an aliquot of a sample extract is measured by adding a first substrate which is the substrate for a first reporter enzyme and a second substrate which is the substrate for a second reporter enzyme to an aliquot of the cell extract. In this
10 preferred embodiment, the first enzyme is a luciferase, the second enzyme is a hydrolytic enzyme, the first substrate is luciferin and the second substrate is a dioxetane. The second reporter enzyme is preferably selected from β -galactosidase, β -glucuronidase, alkaline phosphatase and carboxyl esterase. In a preferred embodiment the second reporter enzyme is β -galactosidase. In this
15 embodiment, the method comprises adding the substrate for the first enzyme at the same time as the substrate for the second enzyme and measuring the chemiluminescent signal by the substrate for the first reporter enzyme by measuring the light signal produced from luciferase-luciferin reaction and simultaneously adding the substrate for the second enzyme. Then the method
20 comprises adding an accelerator solution which substantially inactivates the first reporter enzyme by increasing the pH of the aliquot. The increased pH of the aliquot also activates the light production from the accumulated product of the second reporter enzyme-second substrate reaction. The accelerator solution also includes an enhancer, such as polyvinylbenzyltributyl ammonium chloride, which,
25 as aforesaid, enhances the light signal produced by decomposition of the dioxetane substrate for the second reporter enzyme. The activity of the second reporter enzyme is then measured by reading the chemiluminescent signal intensity in the same aliquot of the cell extract. In embodiments where luciferin is used as the first substrate, the presence of this substrate further enhances the light signal
30 produced by the second substrate.

- 25 -

The method of the present invention also provides a kit for detecting the transcriptional activity of more than one reporter gene in an aliquot of a sample extract. Such a kit comprises the reagents for quantifying each of two or more reporter enzymes and the substrates for each of the reporter enzymes, wherein at least one of the substrates is a dioxetane. The kit also comprises an accelerator solution containing a water soluble polymeric enhancer molecule. Optionally the accelerator solution comprises a water soluble polymeric enhancer molecule at a pH from about 9 to about 14, to raise the pH, decrease the activity of the first reporter enzyme and increase the signal produced from the enzymatic degradation of the second substrate by the second reporter enzyme. The polymeric enhancer preferably comprises bovine serum albumin, human serum albumin or polymeric quaternary onium salts. The polymeric quaternary onium salts preferably comprise polyvinylbenzyltrimethylammonium chloride (TMQ), polyvinylbenzyltributylammonium chloride (TBQ), polyvinylbenzylbenzyltrimethylammonium chloride (BDMQ) or polyvinylbenzyltributylphosphonium chloride.

The following examples are provided to illustrate the present invention and are not intended in any way to limit the scope of the invention.

EXAMPLE I. Preparation of Samples.

Routes

A. Preparation of cell extracts from tissue culture cells:

The lysis solution contained 100 mM potassium phosphate pH 7.8, 0.2% Triton X-100 and stored at 4°C. Dithiothrietol (DTT) was added fresh prior to use to a final concentration of 0.5 mM. The cells were rinsed twice with 1X phosphate buffered saline (PBS) containing: 0.058 Na₂HPO₄, 0.017M NaH₂PO₄, 0.068M NaCl at pH 7.3.

Lysis solution was added to cover cells. For example, 250 μL of lysis buffer was adequate for a 60 mm culture plate. The cells were detached from culture plate using a rubber policeman or equivalent. Non-adherent cells were

- 26 -

pelleted and sufficient lysis buffer added to cover the cells. The cells were then resuspended in the lysis buffer by pipetting.

The lysed cells were then transferred to a microfuge tube and centrifuged
5 for 2 minutes to pellet any debris. The supernatant was then transferred to a fresh microfuge tube. The cell extracts were either used immediately or frozen at -70°C for future use.

B. Preparation of Purified Enzymes

10 Purified luciferase (Analytical Luminescence Laboratories, Ann Arbor, MI) and β -galactosidase (G-5635, Sigma, ST. Louis, MO) were individually diluted in lysis solution (0.1 M potassium phosphate, pH 7.8, 0.2% Triton X-100) containing 0.5 mM DTT 1 mg/ml BSA Fraction V (Sigma A 3059).

15 EXAMPLE II. Chemiluminescent Assay Procedure for Luciferase and β -Galactosidase Reporter Enzymes

All assays were run in triplicate. Buffer A was prepared from lyophilized
pH 7.8 solution containing 75 mM Tricine-Cl pH 7.8, 50 mM MgSO_4 , 7.5 mM
20 ATP, 1.25 mM pyrophosphate, 4 mM EDTA and 1 mg/ml bovine serum albumin (BSA). The lyophilized solution was reconstituted in 5 mL of sterile deionized or Milli-Q water. The lyophilized solution was stored at -20°C before reconstitution.

25 Buffer B was prepared from a lyophilized solution of 0.625 mM luciferin and 5 mM Tris acetate pH 6.5 reconstituted in 20 mL of sterile deionized or Milli-Q water. The β -galactosidase substrate containing 1.34 mg/ml Galacton-Plus™ (100X concentrate, Tropix, Bedford, MA; stored at 4°C or optimally at -20°C) was added to Buffer B, diluted 1:100, immediately before use. Buffer B and the
30 β -galactosidase substrate mixture was preferably used within 24 hours.

2-10 μL of individual cell extracts or purified enzyme were aliquoted into luminometer sample tubes or microplate wells. If less than 10 μL of extract was

- 27 -

used, lysis solution was added to bring total volume up to 10 μ L to ensure the same concentration of reducing agent and other components is present in each assay sample. The amount of cell extract required varied depending on the level of expression and the instrument used. The concentration of extract was adjusted
5 to keep the signal within the linear range of the assay.

Buffer A (25 μ L) was added to the tubes containing the extract samples. Within 10 minutes of adding Buffer A, 100 μ L of Buffer B was injected into the samples. The light signal produced from the luciferase reaction was measured
10 using a 2 second delay and a 5 second integration.

The samples were incubated for 30-60 minutes at room temperature. Then 100 μ L of accelerator was added. The accelerator contained 10% Sapphire II™ (10 mg/ml polyvinyl benzyl tributyl ammonium chloride) and 0.3 M diethanolamine
15 unbuffered at approximately pH 10.8. After a 2 second delay, the light signal produced from the Galacton-Plus™ and β -galactosidase reaction was measured on a monolight 2010 luminometer (Analytical Luminescence Laboratory, Ann Arbor, MI) for 5 seconds.

20 A graph of the relative light units (RLUs) of purified luciferase and β -galactosidase versus the amount of enzyme present showed that 10^{-15} to 10^{-8} g of luciferase and β -galactosidase can be detected by the methods of the present invention.

25 The level of interference from one enzyme on the low level detection of the other enzyme was measured by assaying the individual enzyme dilution series for β -galactosidase and luciferase. Purified luciferase and β -galactosidase were diluted in cell lysis solution. A set of dilutions of each enzyme was assayed following the protocol as described above. Signal intensity following injection of Buffer B into
30 diluted luciferase was depicted graphically. Signal intensity following injection of the accelerator into diluted luciferase was depicted graphically. Signal intensity following injection of accelerator into diluted β -galactosidase was depicted

- 28 -

graphically. The signal intensity following injection of Buffer B into diluted β -galactosidase was depicted graphically. The results indicate that when luciferase is not present, levels of β -galactosidase up to 1 ng do not increase the signal from the luciferase portion of the assay over background. When no β -galactosidase is present, quantities of luciferase below 1 ng do not increase the background signal in the β -galactosidase assay. At luciferase levels 1 ng or above, residual luciferase signal is measureable; however, this level of signal would only interfere with very low level detection of β -galactosidase and a dynamic range over 5 orders of magnitude of enzyme concentration is maintained.

10

EXAMPLE III. Chemiluminescent Assay for Luciferase and β -Glucuronidase Reporter Enzymes

Mammalian cell lysis extracts are prepared as described in Example I. Lysis Buffer A and B, are prepared and stored as described in Example II. The β -glucuronidase substrate (Glucuron™, 100X concentrate, Tropix, Bedford, MA: stored at -20°C) is added to Buffer B, diluted 1:100, immediately before use. Buffer B and the Glucuron™ substrate mixture was preferably used within 24 hours.

20

2-10 μ l of individual cell extracts or purified enzyme diluted in lysis buffer are aliquoted into luminometer sample tubes or microplate wells. If less than 10 μ l of extract is used, lysis solution is added to bring total volume up to 10 μ l to ensure the same concentration of reducing agent and other components is present in each assay sample.

Buffer A (25 μ l) is added to the tubes containing the extract/enzyme samples. Within 10 minutes of adding Buffer A, 100 μ l of Buffer B is injected into the samples. The light signal produced from the luciferase reaction is measured using a 2 second delay and a 5 second measurement.

30

The samples are incubated for 30-60 minutes at room temperature. Then 100 μ l of accelerator (0.3M diethanolamine, 10% Sapphire II™) is added. After a 2

- 29 -

second delay, the light signal produced from the β -glucuronidase reaction is measured for 5 seconds.

5 **EXAMPLE IV. Chemiluminescent Assay Procedure for β -Galactosidase and Alkaline Phosphatase Reporter Enzymes Using Two Dioxetane-Containing Substrates**

10 Cell lysis solution is prepared to contain 0.1 M potassium phosphate, pH 7.8, 0.2 % Triton X-100 as described above. Dithiothriitol (DTT) is added fresh prior to use to a final concentration of 0.5 mM. The cells are rinsed twice with 1X tris buffered saline (TBS) containing : TBS, 0.05 M tris, 0.15 M NaCl, pH7.4. Cell lysis is performed as described above.

15 The cells are then transferred to a microfuge tube and centrifuged for 2 minutes to pellet any debris. The supernatant is then transferred to a fresh microfuge tube. The cell extracts are either used immediately or frozen at -70°C for future use.

20 The cell extract (1-20 μL) is aliquoted into luminometer sample tubes or microplate wells. Reaction buffer (100 μL) containing 0.1 M NaPO_4 , pH 8.0 and containing Galacton™ (0.025mM) is added to the wells. The sample tubes are incubated for 30 minutes at room temperature.

25 100 μL of an accelerator solution containing 0.3 M diethanolamine, 10% Sapphire II™ (polyvinyl benzyl tributyl ammonium chloride, 10% w/w) is injected into the sample tubes and after a 2 second delay, the light signal produced from the β -galactosidase reaction was measured for 5 seconds.

30 After the light signal produced from the Galacton™- β -galactosidase reaction is read, 25-100 μL of CSPD solution was immediately injected into the sample tubes to obtain a final concentration of 0.4 mM CSPD. The samples are incubated for 30 minutes to permit the decay of activated Galacton and for the alkaline

- 30 -

phosphatase/CSPD signal to reach a plateau. The light signal produced by decomposition of CSPD by alkaline phosphatase is measured.

5 Alternatively, the step of incubating the samples for 30 minutes subsequent to the quantification of β -galactosidase is shortened or altogether eliminated as the above assay is performed at higher temperatures. In such a method, the assay is performed at a temperature of 25-40°C.

10 EXAMPLE V. Chemiluminescent Assay for Luciferase, β -Galactosidase and Alkaline Phosphatase

Mammalian cell lysis extracts are prepared as described in Example I. Lysis Buffer and Buffer B are prepared and stored as described in Example II. Buffer A was prepared as in Example II except for the omission of EDTA. The β -galactosidase substrate (Galacton™, 100X concentrate, Tropix, Bedford, MA: stored at -20°C) is added to Buffer B, diluted 1:100, immediately before use. Buffer B and the Galacton™ substrate mixture is preferably used within 24 hours.

20 2-10 μ l of individual cell extracts or purified enzymes diluted in lysis buffer are aliquoted into luminometer sample tubes or microplate wells. If less than 10 μ l of extract is used, lysis solution is added to bring total volume up to 10 μ l to ensure the same concentration of reducing agent and other components is present in each assay sample.

25 Buffer A (25 μ l) is added to the tubes containing the extract/enzyme samples. Within 10 minutes of adding Buffer A, 100 μ l of Buffer B is injected into the samples. The light signal produced from the luciferase reaction is measured using a 2 second delay and 5 second measurement.

30 The samples are incubated for 30-60 minutes at room temperature. Then 100 μ l of accelerator (0.3 M diethanolamine, 10% Sapphire II™) is added. After a 2 second delay, the light signal produced from the β -galactosidase reaction is measured for 5 seconds.

- 31 -

After the light signal produced by the β -galactosidase substrate is read, 25-100 μ l of CSPD solution is immediately injected into the sample tubes to obtain a final concentration of 0.4mM CSPD. The samples are incubated for 30 minutes to permit the decay of activated Galacton™ and for the alkaline phosphatase/CSPD
5 signal to reach a plateau. Following this incubation, the light signal produced by decomposition of CSPD by alkaline phosphatase is measured.

The invention has been described in detail with particular references to the preferred embodiments thereof. However, it will be appreciated that modifications
10 and improvements within the spirit and scope of this invention may be made by those skilled in the art upon considering the present disclosure.

We claim

1. A method of measuring the activity of three or more reporter gene products in an aliquot of a sample extract comprising
5 quantifying the activity of a first reporter enzyme by measuring the light signal produced by degradation of a first substrate by the first reporter enzyme, and
 quantifying the activity of a second reporter enzyme by measuring the light signal produced by the degradation of a second substrate by a second reporter
10 enzyme,
 quantifying the activity of a third reporter enzyme by measuring the light signal produced by the degradation of a third substrate by a third reporter enzyme,
 wherein all quantifications are sequentially performed on the same aliquot of sample extract.
15
2. A method of measuring the activity of at least two reporter gene products in an aliquot of a sample extract comprising
 quantifying the activity of a first reporter enzyme by measuring the light signal produced by degradation of a first substrate by the first reporter enzyme,
20 and
 quantifying the activity of a second reporter enzyme by measuring the light signal produced by the degradation of a second substrate by a second reporter enzyme,
 wherein both quantifications are sequentially performed on the same aliquot
25 of sample extract and wherein the presence of the first substrate enhances the light signal produced by degradation of the second substrate.
3. A method of measuring the activity of at least two reporter gene products in an aliquot of a sample extract comprising
30 quantifying the activity of a first reporter enzyme by measuring the light signal produced by degradation of a first substrate by the first reporter enzyme, and
 and

quantifying the activity of a second reporter enzyme by measuring the light signal produced by the degradation of a second substrate by a second reporter enzyme,

5 wherein both quantifications are sequentially performed on the same aliquot of sample extract and wherein the second substrate is added subsequent to the first substrate.

4. A method of measuring the activity of at least two reporter gene products in an aliquot of a sample extract comprising

10 quantifying the activity of a first reporter enzyme by measuring the light signal produced by degradation of a first substrate by the first reporter enzyme, and

15 quantifying the activity of a second reporter enzyme by measuring the light signal produced by the degradation of a second substrate by a second reporter enzyme,

wherein both quantifications are sequentially performed on the same aliquot of sample extract and wherein the activity of the first reporter enzyme is decreased prior to quantifying the activity of the second reporter enzyme by substantially inactivating the first reporter enzyme by heating the aliquot to degrade the first enzyme or adding inactivating chemicals or substrate analogs or by decreasing the amount of the first substrate.

20

5. The method according to claim 1, 3 or 4, wherein the presence of the first substrate enhances the light signal produced by degradation of the second substrate.

25

6. The method of claim 1, 2 or 4, wherein the first and second substrates are added simultaneously.

7. The method of claim 1, 2 or 4, wherein the second substrate is added subsequent to the first substrate.

30

- 34 -

8. The method according to claim 1, 2 or 3, further comprising decreasing the activity of the first reporter enzyme prior to quantifying the activity of the second reporter enzyme.
- 5 9. The method according to claim 8, wherein decreasing the activity of the first reporter enzyme comprises substantially inactivating the first reporter enzyme or decreasing the amount of the first substrate.
- 10 10. The method according to claim 9, wherein substantially inactivating the first reporter enzyme comprises altering the pH of the aliquot, heating the aliquot to degrade the first enzyme, adding inactivating chemicals or substrate analogs.
- 15 11. The method according to claim 10, wherein substantially inactivating the first reporter enzyme comprises altering the pH of the reaction.
12. The method according to claim 9, wherein decreasing the amount of the first substrate comprises adding an additional amount of the first enzyme sufficient to degrade the first substrate.
- 20 13. The method according to claim 9, wherein decreasing the amount of the first substrate comprises heating the aliquot to degrade the first substrate.
- 25 14. The method according to claim 2 or 3, wherein quantifying the activity of the second reporter enzyme comprises inducing production of light from the degradation of the second substrate by the second reporter enzyme prior to measuring the light signal produced by the degradation of a second substrate by a second reporter enzyme.
- 30 15. The method according to claim 14, wherein the production of light is induced by increasing the pH of the aliquot.

- 35 -

16. The method according to claim 2, 3 or 4, further comprising inducing the activity of the second reporter enzyme by adding the second substrate or activating the second reporter enzyme.
- 5 17. The method according to claim 1, wherein at least two of the first, second or third reporter enzymes are hydrolytic enzymes capable of reacting with a dioxetane.
18. The method according to claim 2, 3 or 4, wherein at least one of the first
10 or second reporter enzymes is a hydrolytic enzyme capable of reacting with a dioxetane.
19. The method according to claim 17 or 18, wherein the first reporter enzyme
15 is selected from the group consisting of luciferase, β -galactosidase, β -glucuronidase, alkaline phosphatase or carboxyl esterase.
20. The method according to claim 19, wherein the first reporter enzyme is luciferase.
- 20 21. The method according to claim 17 or 18, wherein the second reporter enzyme comprises β -galactosidase, β -glucuronidase, alkaline phosphatase or carboxyl esterase.
22. The method according to claim 21, wherein the second reporter enzyme
25 comprises β -galactosidase.
23. The method according to claim 17 wherein the third reporter enzyme
comprises β -galactosidase, β -glucuronidase, alkaline phosphatase or carboxyl
esterase.
30
24. The method according to claim 1, wherein at least one of the first, second or third substrates is a dioxetane.

- 36 -

25. The method according to claim 2, 3 or 4, wherein at least one of the first or second substrates is a dioxetane.
26. The method according to claim 24 or 25, wherein the dioxetane comprises a
5 dioxetane containing a substituted or unsubstituted adamantyl group, and Y group may be substituted or unsubstituted in addition to the enzyme cleavable group.
27. The method according to claim 24 or 25, wherein the dioxetane substrate
10 comprises 3-(4-methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1.1^{3,7}]-
decan]-4-y1-phenyl- β -D-galactopyranoside (Galacton™), and 5-chloro-3-
(methoxyspiro[1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1^{3,7}]-decan-4-y1-phenyl- β -
D-galactopyranoside (Galacton-Plus™).
28. The method according to claim 1, 2, 3 or 4, further comprising adding a
15 water soluble polymeric enhancer molecule to enhance the light signal produced by enzymatic degradation of the dioxetane.
29. The method according to claim 28, wherein the polymeric enhancer
20 comprises bovine serum albumin, human serum albumin or polymeric quaternary onium salts.
30. The method according to claim 29, wherein the polymeric quaternary
onium salts comprise polyvinylbenzyltrimethyl-ammonium chloride (TMQ),
polyvinyl benzyl tributyl ammonium chloride (TBQ) or polyvinylbenzyl
25 benzyldimethylammonium chloride (BDMQ), polyvinylbenzyltributylphosphonium chloride, or polyvinyl tributyl sulfonium chloride.
31. The method according to claim 1, 2, 3 or 4, further comprising adding an
30 accelerator solution prior to quantifying the activity of the second enzyme.
32. The method according to claim 31, wherein the accelerator solution
comprises a water soluble polymeric enhancer molecule at a pH from about 9 to

- 37 -

about 14 which is capable of activating the signal production from the degradation of the second substrate by the second reporter enzyme and inactivate the first reporter enzyme.

- 5 33. The method according to claim 32, wherein the polymeric enhancer comprises bovine serum albumin, human serum albumin or polymeric quaternary onium salts.
34. The method according to claim 33, wherein the polymeric quaternary
10 ammonium salts comprise poly vinylbenzyltrimethyl-ammonium chloride (TMQ), polyvinyl benzyl tributyl ammonium chloride (TBQ) or polyvinylbenzyl benzyldimethylammonium chloride (BDMQ), polyvinylbenzyltributylphosphonium chloride, or polyvinyl tributyl sulfonium chloride.
- 15 35. A method of measuring the activity of more than two reporter gene products in a single aliquot of cell extract comprising
quantifying the activity of a first reporter enzyme in an aliquot of the cell extract by measuring the light signal produced by degradation of a first substrate,
decreasing the activity of the first reporter enzyme,
20 quantifying the activity of a second reporter enzyme in the aliquot of the cell extract by measuring the light signal produced by degradation of a second substrate,
decreasing the activity of the second reporter enzyme, and
quantifying the activity of a third reporter enzyme in the aliquot of the cell
25 extract by measuring the light signal produced by degradation of a third substrate, wherein all quantifications are sequentially performed on the same aliquot of sample extract.
36. The method according to claim 35, wherein the first, second and third
30 substrates are different and at least one of the substrates is a dioxetane.

- 38 -

37. The method according to claim 35, further comprising inducing the signal produced by the degradation of the second substrate by the second reporter enzyme.
- 5 38. The method according to claim 35, further comprising inducing the activity of the third reporter enzyme.
39. A kit for detecting the activity of more than two reporter gene products in an aliquot of a sample extract comprising
- 10 the reagents for quantifying each of more than two reporter enzymes;
the substrates for each of the reporter enzymes;
wherein at least one of the substrates is a dioxetane; and
an accelerator solution containing a water soluble polymeric enhancer molecule.
- 15 40. The kit according to claim 39, wherein the accelerator solution comprises a water soluble polymeric enhancer molecule at a pH from about 9 to about 14, to induce light production from the degradation of the second substrate by the second reporter enzyme and inactivate the first reporter enzyme.
- 20 41. The kit according to claim 39, wherein the polymeric enhancer comprises bovine serum albumin, human serum albumin or polymeric quaternary onium salts.
- 25 42. The kit according to claim 41, wherein the polymeric quaternary ammonium salts comprise poly vinylbenzyltrimethyl-ammonium chloride (TMQ), polyvinyl benzyl tributyl ammonium chloride (TBQ) or polyvinylbenzyl benzyldimethyl ammonium chloride (BDMQ), polyvinylbenzyl-tributylphosphonium chloride, or polyvinyl tributyl sulfonium chloride.
- 30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/20650

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :C12Q 1/68 US CL :435/4, 6, 810 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/4, 6, 810 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, MEDLINE, BIOSIS, CA, DERWENT search terms: reporters, reporter genes, reporter products, luciferase, dioxetane, GUS, B-galactosidase		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y ----- A	BRONSTEIN et al. Chemiluminescent and bioluminescent reporter gene assays. Analytical Biochemistry. 1994, Vol. 219, pages 169-181, especially pages 175-177.	39-42 ----- 1-38
A	ALBERTS et al. Rapid transcriptional assay for the expression of two distinct reporter genes by microinjection. DNA and Cell Biology. 1993, Vol. 12, No. 10, pages 935-943.	1-42
A	US 5,362,865 A (AUSTIN) 08 November 1994, column13, lines 25-50	1-42
A, E	Biotechniques. Vol. 22, No. 1, January 1997, advertisement for Promega product "Dual Luciferase Reporter Assay System".	1-42
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier documents published on or after the international filing date *L* documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family	
Date of the actual completion of the international search 21 FEBRUARY 1997		Date of mailing of the international search report 06 MAR 1997
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer KENNETH R. HORLICK Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/20650

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, E	1997 Promega Catalog, page 229, "Dual-Luciferase Reporter Assay System".	1-42